

NORM-ATTAINING INTEGRAL OPERATORS ON ANALYTIC FUNCTION SPACES

CHENGJI XIONG & JUNMING LIU

ABSTRACT. Any bounded analytic function g induces a bounded integral operator S_g on the Bloch space, the Dirichlet space and $BMOA$ respectively. S_g attains its norm on the Bloch space and $BMOA$ for any g , but does not attain its norm on the Dirichlet space for non-constant g . Some results are also obtained for S_g on the little Bloch space, and for another integral operator T_g from the Dirichlet space to the Bergman space.

1. INTRODUCTION

Given two Banach spaces X and Y , a bounded linear operator T from X to Y is said to be norm-attaining if there is an $x \in X$ with $\|x\| = 1$ such that $\|Tx\| = \|T\|$. Such an x is called an extremal point (or extremal function for the function space X) for the norm of T . Norm-attaining operators have been studied extensively by many authors, such as [10], [7] and [11]. Especially, there are several papers about norm-attaining operators on analytic function spaces. For examples, norm-attaining composition operators on Hardy spaces, Bergman spaces and the Dirichlet space were discussed in [5] and [6] respectively. Recently, María considered norm-attaining composition operators on the Bloch space and the little Bloch space in [9].

Let $\mathbb{D}$ be the open unit disc in the complex plane $\mathbb{C}$ and $H(\mathbb{D})$ the space of all analytic functions on $\mathbb{D}$. Any $g \in H(\mathbb{D})$ induces two integral operators S_g and T_g on $H(\mathbb{D})$ as following

$$S_g f(z) \triangleq \int_0^z f'(w)g(w)dw \quad \text{and} \quad T_g f(z) \triangleq \int_0^z f(w)g'(w)dw.$$

Let M_g denote the multiplication operator $M_g(f) = fg$. Then

$$(S_g + T_g)f = fg - f(0)g(0) = M_g(f) - f(0)g(0).$$

If g is constant, then all results about S_g , T_g or M_g are trivial. In general, g is assumed to be non-constant.

Both integral operators have been studied by several authors. Most recently, bounded below of these integral operators are studied on some analytic functions spaces. See [1] and its references for more materials.

In this paper, the norm-attaining integral operator S_g is investigated on several classic analytic function spaces. In the section 2, the operator norm of S_g on the Bloch space is estimated. We show that S_g is norm-attaining for any g on the Bloch space, but does not attain its norm on the little Bloch space for non-constant

2000 *Mathematics Subject Classification.* 45P05, 30H.

Key words and phrases. integral operator, analytic function space, Norm-attaining.

This work was supported by NNSF of China (Grant No. 11171203).

g . In the section 3, the operator norm of S_g on the Dirichlet space is obtained. We show that S_g does not attain its norm on the Dirichlet space for non-constant g . Another integral operator T_g from the Dirichlet space to the Bergman space, is norm-attaining if and only if g is linear. In the last section, we estimate the operator norm of S_g on $BMOA$ and prove that S_g is norm-attaining on $BMOA$ for any g .

2. Norm-attaining integral operators on $\mathcal{B}_0$ and $\mathcal{B}$

The Bloch space $\mathcal{B}$ is the set of all functions $f \in H(\mathbb{D})$ with

$$\|f\|_{\mathcal{B}} = |f(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| < \infty.$$

The set of all functions in $\mathcal{B}$ satisfying

$$\lim_{|z| \rightarrow 1} (1 - |z|^2) |f'(z)| = 0$$

is called as the little Bloch space $\mathcal{B}_0$. The Bloch space $\mathcal{B}$ becomes a Banach space under the norm $\|\cdot\|_{\mathcal{B}}$ and the little Bloch space is a closed subspace of $\mathcal{B}$. See [14] for more information on Bloch spaces.

A sequence $\{z_n\} \subset \mathbb{D}$ is said to be thin if

$$\lim_{k \rightarrow \infty} \prod_{j, j \neq k} \left| \frac{z_j - z_k}{1 - \overline{z_j} z_k} \right| = 1.$$

It is trivial that a thin sequence implies $z_n \rightarrow \partial\mathbb{D}$. The Blaschke product

$$B(z) = \prod_{n=1}^{\infty} \frac{|z_n|}{z_n} \frac{z_n - z}{1 - \overline{z_n} z}$$

is called thin if its zeros $\{z_n\}$ form a thin sequence. Gorkin and Mortini [4] showed that any sequence in $\mathbb{D}$ tending to the boundary admits a thin subsequence as following.

Lemma A. If $\{z_n\}$ is a sequence in $\mathbb{D}$ and $z_n \rightarrow \partial\mathbb{D}$, then there exists a thin Blaschke product B whose zeros are contained in the set $\{z_n : n \in \mathbb{N}\}$.

Let $B(z)$ be the thin Blaschke product with zeros $\{z_n\}$. Direct computation gives

$$(1 - |z_n|^2) |B'(z_n)| = \prod_{k, k \neq n} \left| \frac{z_n - z_k}{1 - \overline{z_k} z_n} \right|.$$

Using the definition of thin sequence and Schwarz Pick Lemma, it is easy to see

$$1 \geq \sup_{z \in \mathbb{D}} (1 - |z|^2) |B'(z)| \geq \lim_{n \rightarrow \infty} (1 - |z_n|^2) |B'(z_n)| = 1,$$

so the semi-norm of B equals 1.

In [13], Yoneda gave the following result.

Lemma B. A integral operator S_g is bounded on the Bloch space $\mathcal{B}$ ($\mathcal{B}_0$) if and only if the inducing function g belongs to H^∞ , the set of all bounded analytic functions on $\mathbb{D}$.

To determine whether or not an operator is norm-attaining, we need to calculate the operator norm of it. For the operator norm of S_g on $\mathcal{B}$ ($\mathcal{B}_0$), we obtained following result.

Lemma 1. *If a integral operator S_g is bounded on $\mathcal{B}$ ($\mathcal{B}_0$), then*

$$\|S_g\| = \sup_{z \in \mathbb{D}} |g(z)| .$$

Proof. By the Lemma B, we know that g is bounded. For any $f \in \mathcal{B}$ with $\|f\|_{\mathcal{B}} = 1$,

$$\|S_g f\|_{\mathcal{B}} = \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| |g(z)| \leq \|f\|_{\mathcal{B}} \cdot \sup_{z \in \mathbb{D}} |g(z)| = \sup_{z \in \mathbb{D}} |g(z)| ,$$

which implies

$$\|S_g\| \leq \sup_{z \in \mathbb{D}} |g(z)| .$$

Now we need only to show the reverse inequality. Denote $a = \sup_{z \in \mathbb{D}} |g(z)|$. Given any $\epsilon > 0$, there exists $z_0 \in \mathbb{D}$ such that $|g(z_0)| \geq a - \epsilon$. Let

$$f_{z_0}(z) = \frac{z_0 - z}{1 - \bar{z}_0 z} - z_0 .$$

Then $f_{z_0} \in \mathcal{B}_0$, $\|f_{z_0}\|_{\mathcal{B}} = 1$ and $(1 - |z_0|^2) |f'_{z_0}(z_0)| = 1$. We have

$$\begin{aligned} \|S_g\| &\geq \|S_g f_{z_0}\|_{\mathcal{B}} \\ &\geq (1 - |z_0|^2) |f'_{z_0}(z_0)| \cdot |g(z_0)| \\ &\geq a - \epsilon . \end{aligned}$$

Since ϵ is arbitrary, the reverse inequality holds. $\square$

Theorem 1. *If $g \in H^\infty$ is not constant, then the integral operator S_g does not attain its norm on $\mathcal{B}_0$.*

Proof. Suppose that S_g is norm-attaining on $\mathcal{B}_0$, then there exists an extremal function $f \in \mathcal{B}_0$ with $\|f\|_{\mathcal{B}} = 1$ satisfying $\|S_g\| = \|S_g f\|_{\mathcal{B}}$. By Lemma 1, we have

$$\sup_{z \in \mathbb{D}} |g(z)| = \|S_g\| = \|S_g f\|_{\mathcal{B}} = \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| \cdot |g(z)| \leq \sup_{z \in \mathbb{D}} |g(z)| .$$

By definition, f belongs to $\mathcal{B}_0$ implies that

$$\lim_{|z| \rightarrow 1^-} (1 - |z|^2) |f'(z)| = 0 .$$

Hence g must attains its maximal modulus at some point $z_0 \in \mathbb{D}$, which contradicts with Maximal Modulus Theorem since g is not constant by assumption. $\square$

Theorem 2. *For any $g \in H^\infty$, the integral operator S_g attains its norm on Bloch space $\mathcal{B}$. Furthermore, a function $f \in \mathcal{B}$ with $\|f\|_{\mathcal{B}} = 1$ is extremal for the norm of S_g if and only if there exists a sequence $\{z_n\} \subset \mathbb{D}$ with $z_n \rightarrow \partial\mathbb{D}$ such that both conditions*

$$\lim_{n \rightarrow \infty} |g(z_n)| = \sup_{z \in \mathbb{D}} |g(z)| \quad \text{and} \quad \lim_{n \rightarrow \infty} (1 - |z_n|^2) |f'(z_n)| = 1$$

are satisfied.

Proof. If g is constant, it is trivial that S_g is norm-attaining. Given any non-constant $g \in H^\infty$, we will construct an extremal function for the norm of S_g on $\mathcal{B}$. Take a sequence $\{z_n\}$ in $\mathbb{D}$ with $\lim_{n \rightarrow \infty} |g(z_n)| = \sup_{z \in \mathbb{D}} |g(z)|$. Then $\lim_{n \rightarrow \infty} |z_n| = 1$ by Maximal Modulus Theorem. Using Lemma A, we obtain a thin Blaschke product $B(z)$ whose zeros is a subsequence of $\{z_n\}$, which is denoted

also by $\{z_n\}$ for convenience. Let $h(z) = B(z) - B(0)$, then $\|h\|_{\mathcal{B}} = 1$. Keep in mind that

$$\lim_{n \rightarrow \infty} (1 - |z_n|^2) |B'(z_n)| = 1 \text{ and } \lim_{n \rightarrow \infty} |g(z_n)| = \sup_{z \in \mathbb{D}} |g(z)|.$$

We have

$$\begin{aligned} \|S_g\| &\geq \|S_g h\|_{\mathcal{B}} = \sup_{z \in \mathbb{D}} (1 - |z|^2) |B'(z)| \cdot |g(z)| \\ &\geq \lim_{n \rightarrow \infty} (1 - |z_n|^2) |B'(z_n)| \cdot |g(z_n)| \\ &= \sup_{z \in \mathbb{D}} |g(z)| = \|S_g\|, \end{aligned}$$

where the Lemma 1 is used. So the function h is an extremal function for the norm of S_g .

Now, if f is an extremal function for the norm of S_g , then

$$\begin{aligned} \|S_g\| &= \|S_g f\|_{\mathcal{B}} \\ &= \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| \cdot |g(z)| \\ &\leq \sup_{z \in \mathbb{D}} |g(z)| = \|S_g\|. \end{aligned}$$

Since $f \in \mathcal{B}$ and $\|f\|_{\mathcal{B}} = 1$, there exists a sequence $\{z_n\}$ with $z_n \rightarrow \partial\mathbb{D}$ such that

$$\lim_{n \rightarrow \infty} |g(z_n)| = \sup_{z \in \mathbb{D}} |g(z)| \text{ and } \lim_{n \rightarrow \infty} (1 - |z_n|^2) |f'(z_n)| = 1. \quad *$$

Conversely, suppose that there is a sequence $\{z_n\}$ and a unit norm function f such that (*) holds, then

$$\begin{aligned} \|S_g\| &\geq \|S_g f\|_{\mathcal{B}} = \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| \cdot |g(z)| \\ &\geq \lim_{n \rightarrow \infty} (1 - |z_n|^2) |f'(z_n)| \cdot |g(z_n)| \\ &= \sup_{z \in \mathbb{D}} |g(z)| = \|S_g\|. \end{aligned}$$

So f is an extremal function for the norm of S_g . $\square$

3. Norm attaining integral operators on the Dirichlet space

In this section, we discuss norm attaining integral operators on the Dirichlet space. Some related results about another integral operator from the Bergman space to the Dirichlet space are also obtained. At first, let us review definitions of the Dirichlet space and the Bergman space.

Denote $dA(z) = \frac{1}{\pi} dx dy$. The Dirichlet space $\mathcal{D}$ consisting of analytic functions $f \in H(\mathbb{D})$ with

$$\int_{\mathbb{D}} |f'(z)|^2 dA(z) < \infty.$$

The norm of $f \in \mathcal{D}$ is defined as

$$\|f\|_{\mathcal{D}} = \left(|f(0)|^2 + \int_{\mathbb{D}} |f'(z)|^2 dA(z) \right)^{1/2}.$$

The Bergman space $\mathcal{A}^2$ consisting of analytic functions $f \in H(\mathbb{D})$ with

$$\int_{\mathbb{D}} |f(z)|^2 dA(z) < \infty.$$

The norm of $f \in \mathcal{A}^2$ is defined as

$$\|f\|_{\mathcal{A}^2} = \left(\int_{\mathbb{D}} |f(z)|^2 dA(z) \right)^{1/2}.$$

$\mathcal{D}$ and $\mathcal{A}^2$ are Hilbert spaces with

$$\langle f, g \rangle = f(0)\overline{g(0)} + \int_{\mathbb{D}} f'(z)\overline{g'(z)} dA(z)$$

and

$$\langle f, g \rangle = \int_{\mathbb{D}} f(z)\overline{g(z)} dA(z)$$

respectively.

Lemma 2. *Let f be an analytic function in $\mathbb{D}$ and $0 < r < 1$. Then*

$$|f(0)|^2 \leq \frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^2 d\theta \text{ and } |f(0)|^2 \leq \int_{\mathbb{D}} |f(z)|^2 dA(z).$$

Proof. Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ be the Taylor series of f , then

$$|f(z)|^2 = |f(re^{i\theta})|^2 = f(re^{i\theta})\overline{f(re^{i\theta})} = \sum_{j=0, k=0}^{\infty} a_j \overline{a_k} r^{j+k} e^{i(j-k)\theta}.$$

Direct computation by Taylor coefficients gives the result. $\square$

The more general form of above lemma can be found in Zhu's book [14]. In [2], Austin characterize the boundness of integral operators S_g on the Dirichlet space, and obtained following result.

Lemma C. An integral operator S_g is bounded on the Dirichlet space $\mathcal{D}$ if and only if g belongs to H^∞ .

Now we can give the norm of S_g on $\mathcal{D}$.

Lemma 3. *Let S_g be a bounded integral operator on $\mathcal{D}$, then*

$$\|S_g\| = \sup_{z \in \mathbb{D}} |g(z)|.$$

Proof. Since S_g is bounded on $\mathcal{D}$, we can denote $b = \sup_{z \in \mathbb{D}} |g(z)|$ by the Lemma C. For any $\epsilon > 0$, there exists a point $a \in D$ such that $|g(a)| > b - \epsilon$. Consider the function

$$f_a(z) = \sigma_a(z) - a = \frac{a - z}{1 - \overline{a}z} - a,$$

then $\|f_a\|_{\mathcal{D}} = 1$. By the Lemma 2, we have

$$\begin{aligned} \|S_g\| &\geq \|S_g f_a\|_{\mathcal{D}} \\ &= \left(\int_{\mathbb{D}} |g(z)|^2 |\sigma'_a(z)|^2 dA(z) \right)^{1/2} \\ &= \left(\int_{\mathbb{D}} |g(\sigma_a(z))|^2 dA(z) \right)^{1/2} \\ &\geq |g(\sigma_a(0))| = |g(a)| > b - \epsilon. \end{aligned}$$

Hence $\|S_g\| \geq \sup_{z \in \mathbb{D}} |g(z)|$.

The reverse inequality can be obtained by

$$\|S_g f\|_{\mathcal{D}} = \left(\int_{\mathbb{D}} |f'(z)g(z)|^2 dA(z) \right)^{1/2} \leq \|f\|_{\mathcal{D}} \cdot \sup_{z \in \mathbb{D}} |g(z)|$$

for any $f \in \mathcal{D}$. □

Using above Lemma 3, we obtain main result of this section.

Theorem 3. *Let g be a non-constant function in H^∞ . Then the integral operator S_g can not attain its norm on the Dirichlet space $\mathcal{D}$.*

Proof. Assume that S_g attains its norm and f is an extremal function. Then it is necessary that $f(0) = 0$. Otherwise, we have

$$\begin{aligned} \|S_g\| &= \|S_g f\|_{\mathcal{D}} \\ &= \left(\int_{\mathbb{D}} |f'(z)|^2 |g(z)|^2 dA(z) \right)^{1/2} \\ &\leq \left(\int_{\mathbb{D}} |f'(z)|^2 dA(z) \right)^{1/2} \cdot \sup_{z \in \mathbb{D}} |g(z)| \\ &< \left(|f(0)|^2 + \int_{\mathbb{D}} |f'(z)|^2 dA(z) \right)^{1/2} \cdot \sup_{z \in \mathbb{D}} |g(z)| \\ &= \sup_{z \in \mathbb{D}} |g(z)| = \|S_g\|, \end{aligned}$$

where $\|f\|_{\mathcal{D}} = 1$ and the Lemma 3 is used. Then

$$\int_{\mathbb{D}} |f'(z)|^2 dA(z) = 1$$

and

$$\|S_g\|^2 = \|S_g f\|_{\mathcal{D}}^2 = \int_{\mathbb{D}} |f'(z)|^2 |g(z)|^2 dA(z).$$

But this is impossible since

$$\int_{\mathbb{D}} |f'(z)|^2 (|g(z)|^2 - \|S_g\|^2) dA(z) < 0$$

for any non-constant analytic function g . □

Consider another integral operator $T_g f(z) = \int_0^z g'(w) f(w) dw$. Following results can be deduced similarly with the Lemma 3 and the Theorem 3. We omit proofs of them.

Lemma 4. *If the integral operator T_g is bounded from A^2 to $\mathcal{D}$, then*

$$\|T_g\| = \sup_{z \in \mathbb{D}} |g'(z)|.$$

This lemma can be proved by replacing f_a with

$$F_a(z) = \frac{1 - |a|^2}{(1 - \bar{a}z)^2}, \quad a \in \mathbb{D}$$

in the proof of the Lemma 3.

Theorem 4. *Let T_g be a bounded operator from A^2 to $\mathcal{D}$. Then T_g is norm attaining if and only if $g(z) = az + b$, where $a, b \in \mathbb{C}$.*

4. Norm attaining integral operators on $BMOA$

The analytic Hardy space H^2 on the unit disc $\mathbb{D}$ consists of all analytic functions $f \in H(\mathbb{D})$ satisfying

$$\|f\|_{H^2} = \left(\sup_{0 < r < 1} \frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^2 d\theta \right)^{1/2} < \infty .$$

For $a \in \mathbb{D}$, denote

$$\sigma_a(z) = \frac{a - z}{1 - \bar{a}z}, \quad z \in \mathbb{D} .$$

Let $BMOA$ denote the space of all analytic functions $f \in H^2$ whose boundary function have bounded mean oscillation. The norm of $f \in BMOA$ can be defined as

$$\|f\|_{BMOA} = |f(0)| + \sup_{a \in \mathbb{D}} \|f \circ \sigma_a - f(a)\|_{H^2} .$$

There are some other equivalent norms for $BMOA$, see [3], for example. Next is the well-known Littlewood-Paley Identity, whose proof can be found in many books.

Lemma D. Let f be analytic on unit disc $\mathbb{D}$. Then

$$\|f\|_{H^2}^2 = |f(0)|^2 + 2 \int_{\mathbb{D}} |f'(z)|^2 \log \frac{1}{|z|} dA(z) .$$

By Littlewood-Paley Identity, the norm of $BMOA$ can be expressed as

$$\|f\|_{BMOA} = |f(0)| + \sup_{a \in \mathbb{D}} \left\{ \int_{\mathbb{D}} 2|f'(z)|^2 \log \frac{1}{|\sigma_a(z)|} dA(z) \right\}^{1/2} .$$

The following lemma is our main tool to compute the norm of integral operators.

Lemma 5. If f is analytic in $\mathbb{D}$, then

$$|f(0)|^2 \leq 2 \int_{\mathbb{D}} |f(z)|^2 \log \frac{1}{|z|} dA(z) .$$

Proof. The proof is similar to that of the Lemma 2. □

In [12], Xiao showed that S_g is bounded on $BMOA$ if and only if $g \in H^\infty$. Now we give the norm of S_g on $BMOA$.

Lemma 6. If S_g is a bounded integral operator on $BMOA$, then

$$\|S_g\| = \sup_{z \in \mathbb{D}} |g(z)| .$$

Proof. Since S_g is bounded on $BMOA$, g is bounded by Xiao [12]. Let $\lambda = \sup_{z \in \mathbb{D}} |g(z)|$. For any $\epsilon > 0$, there exists a point $b \in \mathbb{D}$ such that $|g(b)| > \lambda - \epsilon$.

Denote $f_b(z) = \sigma_b(z) - b$. Then

$$\begin{aligned}
\|f_b\|_{BMOA} &= \sup_{a \in \mathbb{D}} \left\{ \int_{\mathbb{D}} 2|\sigma'_b(z)|^2 \log \frac{1}{|\sigma_a(z)|} dA(z) \right\}^{1/2} \\
&= \sup_{a \in \mathbb{D}} \left\{ \int_{\mathbb{D}} 2|[\sigma_b(\sigma_a(z))]'|^2 \log \frac{1}{|z|} dA(z) \right\}^{1/2} \\
&= \sup_{c \in \mathbb{D}} \left\{ \int_{\mathbb{D}} 2|\sigma'_c(z)|^2 \log \frac{1}{|z|} dA(z) \right\}^{1/2} \\
&= \sup_{c \in \mathbb{D}} \left\{ (\|\sigma_c\|_{H^2}^2 - |c|^2) \right\}^{1/2} \\
&= \sup_{c \in \mathbb{D}} (1 - |c|^2)^{1/2} = 1
\end{aligned}$$

by the Lemma D. Hence

$$\begin{aligned}
\|S_g\| &\geq \|S_g f_b\|_{BMOA} \\
&= \sup_{a \in \mathbb{D}} \left(\int_{\mathbb{D}} 2|f'_b(z)g(z)|^2 \log \frac{1}{|\sigma_a(z)|} dA(z) \right)^{1/2} \\
&= \sup_{a \in \mathbb{D}} \left(\int_{\mathbb{D}} 2|g(z)|^2 |\sigma'_b(z)|^2 \log \frac{1}{|\sigma_a(z)|} dA(z) \right)^{1/2} \\
&= \sup_{a \in \mathbb{D}} \left(\int_{\mathbb{D}} 2|g(\sigma_b(z))|^2 \log \frac{1}{|\sigma_a(\sigma_b(z))|} dA(z) \right)^{1/2} \\
&\geq \left(2 \int_{\mathbb{D}} |g(\sigma_b(z))|^2 \log \frac{1}{|z|} dA(z) \right)^{1/2} \\
&\geq |g(b)| \geq \lambda - \epsilon,
\end{aligned}$$

where the Lemma 5 is used.

The reverse inequality can be followed from

$$\|S_g f\|_{BMOA} = \sup_{a \in \mathbb{D}} \left(\int_{\mathbb{D}} 2|f'g|^2 g(z, a) dA(z) \right)^{1/2} \leq \|f\|_{BMOA} \cdot \sup_{z \in \mathbb{D}} |g(z)|.$$

The proof is completed. $\square$

Next theorem is the main result of this section.

Theorem 5. *For any $g \in H^\infty$, the integral operator S_g is norm-attaining on $BMOA$.*

Proof. Similar to the proof of the Theorem 2, we need only to construct an extremal function for non-constant g . Choose a sequence $\{z_n\} \subset \mathbb{D}$ with $\lim_{n \rightarrow \infty} |g(z_n)| = \sup_{z \in \mathbb{D}} |g(z)|$ since $g \in H^\infty$. Using the Lemma A, we obtain a thin Blaschke product $B(z)$ whose zeros is a subsequence of $\{z_n\}$, which is denoted also by $\{z_n\}$ for convenience. Consider $h(z) = B(z) - B(0)$, then $\|h\|_{BMOA} = 1$ by Proposition 2.2 of [8]. Note that

$$\lim_{n \rightarrow \infty} (1 - |z_n|^2) |h'(z_n)| = 1 \quad \text{and} \quad \lim_{n \rightarrow \infty} |g(z_n)| = \sup_{z \in \mathbb{D}} |g(z)|.$$

We have

$$\begin{aligned}
\|S_g h\|_{BMOA} &= \sup_{a \in \mathbb{D}} \left(\int_{\mathbb{D}} 2|h'(z)|^2 |g(z)|^2 \log \frac{1}{|\sigma_a(z)|} dA(z) \right)^{1/2} \\
&= \sup_{a \in \mathbb{D}} \left(2 \int_{\mathbb{D}} |h'(\sigma_a(w))| |g(\sigma_a(w))|^2 |\sigma'_a(w)|^2 \log \frac{1}{|w|} dA(w) \right)^{1/2} \\
&\geq \left(2 \int_{\mathbb{D}} |h'(\sigma_{z_n}(w))| |g(\sigma_{z_n}(w))|^2 |\sigma'_{z_n}(w)|^2 \log \frac{1}{|w|} dA(w) \right)^{1/2} \\
&\geq (1 - |z_n|^2) |h'(z_n)| |g(z_n)|,
\end{aligned}$$

where the Lemma 5 is used. Let $n \rightarrow \infty$, then $\|S_g h\|_{BMOA} \geq \sup_{z \in \mathbb{D}} |g(z)|$ and $\|S_g h\|_{BMOA} = \sup_{z \in \mathbb{D}} |g(z)| = \|S_g\|$ since $\|h\|_{BMOA} = 1$. Hence the function h is an extremal function for the norm of S_g and S_g is norm-attaining on $BMOA$. $\square$

REFERENCES

- [1] Austin, A, *Some closed range integral operators on spaces of analytic functions*, Integral Equations Operator Theory 69 (2011), no. 1, 87-99.
- [2] Austin, A, *Multiplication and Integral Operators on Banach Spaces of Analytic Functions*, PhD Thesis, University of Hawaii, 2010.
- [3] Girela, Daniel, *Analytic functions of bounded mean oscillation*, Complex function spaces (Mekrijärvi, 1999), 61-170, Univ. Joensuu Dept. Math. Rep. Ser., 4, Univ. Joensuu, Joensuu, 2001.
- [4] Gorkin, P. and Mortini, R., *Universal Blaschke products*, Math. Proc. Cambridge Philos. Soc. 136 (2004), no. 1, 175-184.
- [5] Hammond, Christopher, *On the norm of a composition operator with linear fractional symbol*, Acta Sci. Math. (Szeged) 69 (2003), no. 3-4, 813-829.
- [6] Hammond, Christopher, *The norm of a composition operator with linear symbol acting on the Dirichlet space*, J. Math. Anal. Appl. 303 (2005), no. 2, 499-508.
- [7] Khatskevich, V. A.; Ostrovskii, M. I. and Shulman, V. S., *Extremal problems for operators in Banach spaces arising in the study of linear operator pencils*, Integral Equations Operator Theory 51 (2005), no. 1, 109-119.
- [8] Laitila, J., *Isometric composition operators on BMOA*, Math. Nachr. 283 (2010), no. 11, 1646-1653.
- [9] Martía, M. J., *Norm-attaining composition operators on the Bloch spaces*, J. Math. Anal. Appl. 369 (2010) no. 1, 15-21.
- [10] Schachermayer, W., *Norm attaining operators on some classical Banach spaces*, Pacific J. Math. 105 (1983), no. 2, 427-438.
- [11] Shkarin, S., *Norm attaining operators and pseudospectrum*, Integral Equations Operator Theory 64 (2009), no. 1, 115-136.
- [12] Xiao, J., *The Q_p Carleson measure problem*, Adv. Math. 217 (2008), no. 5, 2075-2088.
- [13] Yoneda, R., *Multiplication operators, integration operators and companion operators on weighted Bloch space*, Hokkaido Math. J. 34 (2005), no. 1, 135-147.
- [14] Zhu, K., *Operator Theory in Function Spaces*, Second edition. Mathematical Surveys and Monographs, 138. Amer. Math. Soc, Providence (2007).

SCHOOL OF MATHEMATICAL SCIENCE, HUAIYIN NORMAL UNIVERSITY, HUAIAN, JIANGSU, 223300,
P. R. CHINA
E-mail address: chengji_xiong@yahoo.com

DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG, 515063,
P. R. CHINA
E-mail address: 08jmliu@stu.edu.cn